

*Afr J Tradit Complement Altern Med*. 2011; 8(5 Suppl): 226–232.

PMCID: PMC3252704

Published online 2011 Jul 3. doi: [10.4314/ajtcam.v8i5S.10](https://doi.org/10.4314/ajtcam.v8i5S.10)

PMID: [22754079](https://pubmed.ncbi.nlm.nih.gov/22754079/)

## Anticancer Activities of *Nigella Sativa* (Black Cumin)

Md Asaduzzaman Khan,<sup>1</sup> Han-chun Chen,<sup>✉1</sup> Mousumi Tania,<sup>1</sup> and Dian-zheng Zhang<sup>1,2</sup>

<sup>1</sup>Department of Biochemistry, School of Biological Science and Technology, Central South University, Changsha, Hunan 410013, P R China

<sup>2</sup>Department of Biochemistry/Molecular Biology, Philadelphia College of Osteopathic Medicine, Philadelphia, PA 19131, USA

✉Corresponding author.

\*E-mail: [asadkhanbmj@yahoo.com](mailto:asadkhanbmj@yahoo.com), [chenhanchun@mail.csu.edu.cn](mailto:chenhanchun@mail.csu.edu.cn)

Copyright © Afr. J. Traditional Complementary and Alternative Medicines 2011

### Abstract

*Nigella sativa* has been used as traditional medicine for centuries. The crude oil and thymoquinone (TQ) extracted from its seeds and oil are effective against many diseases like cancer, cardiovascular complications, diabetes, asthma, kidney disease etc. It is effective against cancer in blood system, lung, kidney, liver, prostate, breast, cervix, skin with much safety. The molecular mechanisms behind its anticancer role is still not clearly understood, however, some studies showed that TQ has antioxidant role and improves body's defense system, induces apoptosis and controls Akt pathway. Although the anti-cancer activity of *N. sativa* components was recognized thousands of years ago but proper scientific research with this important traditional medicine is a history of last 2~3 decades. There are not so many research works done with this important traditional medicine and very few reports exist in the scientific database. In this article, we have summarized the actions of TQ and crude oil of *N. sativa* against different cancers with their molecular mechanisms.

**Keywords:** Traditional medicine, *Nigella sativa*, Thymoquinone, Antioxidant, Anti-cancer mechanism


### Introduction

Cancer is one of the major threats of modern life, which is considered as the second cause of death after myocardial infarction ([Grundy, 1991](#)). Millions of people die every year in different types of cancer despite tremendous efforts to find methods of control and cure. In the last century, great advances were made in modern medical science to control disease. But many diseases like cancers are not yet curable fully. To find out new and authentic therapies, scientists are working with traditional or folk medicines in parallel of modern medicine. *Nigella sativa* has been used for medicinal purposes for centuries. It originated from Southeastern Asia and also used in ancient Egypt, Greece, Middle East and Africa. In Islam, it is regarded as one of the greatest forms of healing medicine available ([Nigella-sativa-research.com, 2010](#); [Wikipedia, 2010](#)). It is a flowering plant, of which seed is used as a spice. The seed is called black cumin in English, while in old Latin it was called 'Panacea' meaning 'cure all'; in Arabic it is termed as 'Habba Sawda' or 'Habbat el Baraka' translated as 'Seeds of blessing'. It is also known as 'Kalo jeera' (in Bangladesh), 'Kalonji' (in India) and 'Hak Jung Chou' in (China) ([Aggarwal et al., 2008](#)). Both seeds and oil extracted from this plant are used in medicinal purposes.


The active ingredients of *N. sativa* have beneficial effects against many diseases, including cancers. For example, it is effective in the diminishing the risk of atherosclerosis by decreasing the serum low density lipoprotein cholesterol level and increasing the serum high density lipoprotein cholesterol levels ([Dahri et al., 2005](#); [Nader et al., 2010](#)); it exerts therapeutic and protective effect in diabetes by decreasing morphological changes and preserving pancreatic beta-cell integrity ([Kanter et al., 2009](#)) and by beneficially changing the hepatic enzyme activities ([Pari and Sankaranarayanan, 2009](#)); it is effective against hypertension ([Khattab and Nagi, 2007](#); [Dehkordi and Kamkhah, 2008](#)); it has a potent antihistaminic effect on airways of asthmatic patients ([Boskabady et al., 2010](#)); its components are promising agents to complement schistosomiasis specific treatment ([El Shenawy et al., 2008](#)); its oil protects kidney tissue against oxygen free radicals, preventing renal dysfunction and morphological abnormalities ([Bayrak et al., 2008](#); [Uz et al., 2008](#); [Ragheb et al., 2009](#)). For thousands of year, the seeds, oils and extracts of *N. sativa* have been used as an anticancer agent by Unani, Ayurveda and the Chinese system of medicine that have originated from the Arab, Ind-Bangla and China, respectively. The modern scientific research with the investigation of anticancer activity of *N. sativa* is a comparatively recent affair (for the last 2~3 decades). There are not so many research works done in this field and very few review articles exist in this area. We have searched the scientific databases like Pubmed, Web of Science and Google scholar and summarized the current scientific information about the anticancer activities of *N. sativa* with mechanisms of action.

### Role of *N. sativa* as an anticancer agent

Many active ingredients have been found in the seeds of *N. sativa*. The seeds contain both fixed and essential oils, proteins, alkaloids and saponin ([Ali and Blunden, 2003](#)). [Ghosheh et al. \(1999\)](#) described the quantification of four pharmacologically important components: thymoquinone (TQ), dithymoquinone (DTQ), thymohydroquinone (THQ), and thymol (THY), in the oil of *N. sativa* seed by HPLC. Much of the biological activities of the seeds have been shown to be due to thymoquinone, the major component of the essential oil, which is also present in the fixed oil ([Ali and Blunden, 2003](#)). TQ is considered as potent anti-oxidant ([Badary et al., 2003](#)), anti-carcinogenic and anti-mutagenic agent ([Bourgou et al., 2008](#); [Khader et al., 2010](#)) (structure of thymoquinone is shown in [Figure 1a](#)). Moreover, TQ is a relatively safe compound, particularly when given orally to experimental animals ([Al-Ali et al., 2008](#)). Alpha ( $\alpha$ )-hederin, a pentacyclic triterpene saponin (structure: [Figure 1b](#)) isolated from the seeds of *N. sativa*, was also reported to have potent *in vivo* antitumor activity ([Swamy and Huat, 2003](#)).



(a) Thymoquinone ((source: PubChem-CID [10281](#))



[Open in a separate window](#)

Figure 1

Chemical structure of anti-tumor agents isolated from *N. sativa*

*N. sativa* seeds or oils or its active ingredients like TQ are effective against different cancers:

### Blood Cancer

[El-Mahdy et al. \(2005\)](#) reported that TQ exhibits anti-proliferative effect in human myeloblastic leukemia HL-60 cells. Derivatives of TQ bearing terpene-terminated 6-alkyl residues were tested in HL-60 cells and 518A2 melanoma by [Effenberger et al. \(2010\)](#). They found the derivatives induce apoptosis associated with DNA laddering, a decrease in mitochondrial membrane potential and a slight increase in reactive oxygen species. [Swamy and Huat \(2003\)](#) observed that  $\alpha$ -hederin also induced death of murine leukemia P388 cells by a dose- and time-dependent increase in apoptosis.

### Breast Cancer

---

Aqueous and alcohol extracts of *N. sativa* were found to be effective *in vitro* in inactivating MCF-7 breast cancer cells ([Farah and Begum, 2003](#)). *N. sativa*, in combination with melatonin and retinoic acid reduced the carcinogenic effects of DMBA (7, 12-di-methylbenz(a)anthracene) in mammary carcinoma of rats ([El-Aziz et al., 2005](#)). Terpene-terminated 6-alkyl residues of TQ were tested in MCF-7/Topo breast carcinoma by [Effenberger et al. \(2010\)](#). They found the derivatives inducing cell death by apoptosis.

### Colon Cancer

---

[Gali-Muhtasib et al. \(2004\)](#) suggested that TQ is anti-neoplastic and pro-apoptotic against colon cancer cell line HCT116. [Salim and Fukushima \(2003\)](#) demonstrated that the volatile oil of *N. sativa* has the ability to inhibit colon carcinogenesis of rats in the post-initiation stage, with no evident adverse side effects. [Norwood et al. \(2006\)](#) suggested TQ as chemotherapeutic agent on SW-626 colon cancer cells, in potency, which is similar to 5-fluorouracil in action. However, on HT-29 (colon adenocarcinoma) cell, no effect of TQ was found ([Rooney and Ryan, 2005](#)).

### Pancreatic Cancer

---

[Chehl et al. \(2009\)](#) showed that TQ, the major constituent of *N. sativa* oil extract, induced apoptosis and inhibited proliferation in PDA (pancreatic ductal adenocarcinoma) cells. They also suggested TQ as a novel inhibitor of pro-inflammatory pathways, which provides a promising strategy that combines anti-inflammatory and proapoptotic modes of action. TQ also can abrogate gemcitabine- or oxaliplatin-induced activation of NF-kappa B, resulting in the chemosensitization of pancreatic tumors to conventional therapeutics ([Banerjee et al., 2009](#)). The high molecular weight glycoprotein mucin 4 (MUC4) is aberrantly expressed in pancreatic cancer and contributes to the regulation of differentiation, proliferation, metastasis, and the chemoresistance of pancreatic cancer cells. [Torres et al. \(2010\)](#) evaluated the down-regulatory effect of TQ on MUC4 in pancreatic cancer cells. But in a study, [Rooney and Ryan \(2005\)](#) did not find any preventive role of TQ on MIA PaCa-2 (pancreas carcinoma) cells.

### Hepatic Cancer

---

The cytotoxic activity of *N. sativa* seed was tested on the human hepatoma HepG2 cell line by [Thabrew et al. \(2005\)](#), and 88% inhibitory effect on HepG2 was found after 24-hr incubation with different concentrations (0–50 mg/ml) of the *N. sativa* extract. [Nagi and Almakki \(2009\)](#) reported that oral administration of TQ is effective in increasing the activities of quinone reductase and glutathione transferase and makes TQ a promising prophylactic agent against chemical carcinogenesis and toxicity in hepatic cancer

### Lung Cancer

---

[Swamy and Huat \(2003\)](#) mentioned the antitumor activity of  $\alpha$ -hederin from *N. sativa* against LL/2 (Lewis Lung carcinoma) in BDF1 mice. Also, [Mabrouk et al. \(2002\)](#) showed that supplementation of diet with honey and *N. sativa* has a protective effect against MNU (methylnitrosourea)-induced

oxidative stress, inflammatory response and carcinogenesis in lung, skin and colon. However, [Rooney and Ryan \(2005\)](#) reported that  $\alpha$ -hederin and TQ, the two principal bioactive constituents of *N. sativa* enhance neither cytotoxicity nor apoptosis in A549 (lung carcinoma), HEP-2 (larynx epidermoid carcinoma) cells.

### Skin cancer

---

Topical application of *N. sativa* extract inhibited two-stage initiation/promotion [dimethylbenz[a]anthracene (DMBA)/croton oil] skin carcinogenesis in mice. Again, intraperitoneal administration of *N. sativa* (100 mg/kg body wt) 30 days after subcutaneous administration of MCA (20-methylcholanthrene) restricted soft tissue sarcomas to 33.3% compared with 100% in MCA-treated controls ([Salomi et al., 1991](#)).

### Fibrosarcoma

---

TQ from *N. sativa* was administrated (0.01% in drinking water) one week before and after MCA treatment significantly inhibited the tumor incidence (fibrosarcoma) and tumor burden by 43% and 34%, respectively, compared with the results in the group receiving MCA alone. Moreover, TQ delayed the onset of MCA-induced fibrosarcoma tumors. Also *in vitro* studies showed that TQ inhibited the survival of fibrosarcoma cells with IC<sub>50</sub> of 15 mM ([Badary and Gamal, 2001](#)). Oil of *N. sativa* also decreased the fibrinolytic potential of the human fibrosarcoma cell line (HT1080) *in vitro* ([Awad, 2005](#)).

### Renal Cancer

---

[Khan and Sultana \(2005\)](#) reported the chemo-preventive effect of *N. sativa* against ferric nitrilotriacetate (Fe-NTA)-induced renal oxidative stress, hyper-proliferative response and renal carcinogenesis. Treatment of rats orally with *N. sativa* (50 100 mg/kg body wt) resulted in significant decrease in H<sub>2</sub>O<sub>2</sub> generation, DNA synthesis and incidence of tumors.

### Prostate Cancer

---

TQ, from *N. sativa*, inhibited DNA synthesis, proliferation, and viability of cancerous (LNCaP, C4-B, DU145, and PC-3) but not non-cancerous (BPH-1) prostate epithelial cells by down-regulating AR (androgen receptor) and E2F-1 (a transcription factor) ([Kaseb et al., 2007](#)). In this study, they suggested TQ as effective in treating hormone-sensitive as well as hormone-refractory prostate cancer. [Yi et al. \(2008\)](#) found that TQ blocked angiogenesis *in vitro* and *in vivo*, prevented tumor angiogenesis in a xenograft human prostate cancer (PC3) model in mouse, and inhibited human prostate tumor growth at low dosage with almost no chemotoxic side effects. Furthermore, they observed that endothelial cells were more sensitive to TQ-induced cell apoptosis, cell proliferation, and migration inhibition compared with PC3 cancer cells. TQ also inhibited vascular endothelial growth factor-induced extracellular signal-regulated kinase activation but showed no inhibitory effects on vascular endothelial growth factor receptor 2 activation.

### Cervical Cancer


---

[Shafi et al. \(2009\)](#) reported that methanol, n-Hexane and chloroform extracts of *N. sativa* effectively killed HeLa (human epithelial cervical cancer) cells by inducing apoptosis. [Effenberger et al. \(2010\)](#) tested terpene-terminated 6-alkyl residues of TQ on multidrug-resistant KB-V1/Vb1 cervical carcinoma and found the derivatives inducing cell death by apoptosis.

### Molecular mechanisms of *N. sativa* action against cancer

---

Cancers are the abnormal cell growth caused by genetic alteration. So, any agent which has anti-cancer activity, either protect genetic material from alteration or kill the genetically altered cancer cells. The active ingredients (mainly TQ) from *N. sativa* act on cancer cell to help to kill them by several molecular pathways (Figure 2).



**Figure 2**

Possible mechanisms of thymoquinone (TQ) action. (1) TQ induces apoptotic cell death in cancerous tissues by up-regulating expression of apoptotic genes (caspases and bax) and down-regulating expression of anti-apoptotic genes (e.g., bcl 2); (2) TQ suppresses Akt activation by dephosphorylation and thus blocks cancer cell survival; (3) TQ deactivates NF-kappa B pathway by inducing cytokine production, and thus control oncogenic expression; (4) TQ increases the activities of antioxidant enzymes and protects cell against cancer; (5) TQ protects normal cells' injury caused by ionizing radiation in the treatment of cancer; (6) TQ prevents CYP450 enzymes from damage. '+' indicates increasing effect and '-' indicates decreasing effect.

El-Mahdy et al. (2005) suggested the apoptotic mechanisms behind the anti-proliferative effect of TQ (from *N. sativa*) on myeloblastic leukemia HL-60 cells. They reported that TQ induces apoptosis, disrupts mitochondrial membrane potential and triggers the activation of caspases 8, 9 and 3 in HL-60 cells. The apoptosis induced by TQ was inhibited by a general caspase inhibitor, z-VAD-FMK; a caspase-3-specific inhibitor, z-DEVD-FMK; as well as a caspase-8-specific inhibitor, z-IETD-FMK. Moreover, the caspase-8 inhibitor blocked the TQ-induced activation of caspase-3, PARP cleavage and the release of cytochrome c from mitochondria into the cytoplasm. In addition, TQ treatment of HL-60 cells caused a marked increase in Bax/Bcl2 ratios due to upregulation of Bax and downregulation of

Bcl2 proteins. Their results indicated that TQ-induced apoptosis is associated with the activation of caspases 8, 9 and 3, with caspase-8 acting as an upstream activator and activated caspase-8 initiates the release of cytochrome c during TQ-induced apoptosis. TQ action was also found as pro-apoptotic against colon cancer cell line HCT116 ([Gali-Muhtasib et al., 2004](#)). It was showed that the apoptotic effects of TQ are modulated by Bcl-2 protein and are linked to and dependent on p53. TQ also down-regulates the expression of NF-kappa B-regulated antiapoptotic (IAP1, IAP2, XIAP Bcl-2, Bcl-xL, and survivin) gene products ([Sethi et al., 2008](#)). [Torres et al. \(2010\)](#) found TQ inducing apoptosis by the activation of c-Jun NH(2)-terminal kinase and p38 mitogen-activated protein kinase pathways in pancreatic cancer cell.

TQ has also been reported to be active in controlling Akt pathway. [Yi et al. \(2008\)](#) found that TQ effectively inhibited human umbilical vein endothelial cell migration, invasion, and tube formation by suppressing the activation of AKT and extracellular signal-regulated kinase. [Xuan et al. \(2010\)](#) found that LPS (lipopolysaccharides: a bacterial component)-induced phosphorylation of prosurvival kinases Akt and ERK1/2 was abrogated by TQ in dendritic cells.

NF-kappa B plays a key role in regulating the immune response, and incorrect regulation of NF-kappa B has been found to be linked to cancer ([Albensi and Mattson, 2000](#)). [Sethi et al. \(2008\)](#) found that TQ suppressed tumor necrosis factor-induced NF-kappa B activation in a dose- and time-dependent manner and inhibited NF-kappa B activation induced by various carcinogens and inflammatory stimuli. The suppression of NF-kappa B activation is correlated with sequential inhibition of the activation of I kappa B alpha kinase, I kappa B alpha phosphorylation, I-kappa-B-alpha degradation, p65 phosphorylation, p65 nuclear translocation, and the NF-kappa B-dependent reporter gene expression. Also [Oberge et al. \(2009\)](#) reported that a herbal melanin (HM) from *N. sativa* modulates cytokine production and suggested it as a ligand for TLR4 (toll-like receptor 4). They investigated the possibility that the HM-induced cytokine production is via an NF-kappa B signaling pathway and found that HM induced the degradation of I kappa B-alpha, a key step in the activation of NF-kappa B. Moreover, addition of I kappa B kinase (IKK) specific inhibitors effectively inhibited the observed HM-induced production of IL-8 and IL-6 by TLR4-transfected HEK293 (embryonic kidney 293) cells and THP-1 (Human acute monocytic leukemia) cells ([Oberge et al., 2009](#)).

Many studies showed that *N. sativa* oil or TQ has antioxidant activity and increases the activities of antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT) glutathione peroxidase (GPx) etc ([Ebru et al., 2008](#); [Barron et al., 2008](#); [Ismail et al., 2010](#)). And antioxidant enzymes are clearly related to cancer- mostly their increased activities are beneficial against different types of cancer ([Khan et al., 2010](#)). Administration of *N. sativa* oil or TQ can lower the toxicity of other anticancer drugs (for example, cyclophosphamide) by an up-regulation of antioxidant mechanisms, indicating a potential clinical application for these agents to minimize the toxic effects of treatment with anticancer drugs ([Alenzi et al., 2010](#)).

In addition to these cancer inhibiting properties, components of *N. sativa* have cancer protective roles. [Ibrahim et al. \(2008\)](#) reported that *N. sativa* oil administration has a protective effect against the CCl<sub>4</sub>-mediated suppression of CYP (drug-metabolizing cytochrome P450 enzymes). And genetic abnormalities and polymorphisms of CYP enzymes are associated with cancer ([Sim and Ingelman-Sundberg, 2006](#); [Chen et al., 2008](#)). Radiotherapy is one of the most common strategies for treating human cancers but this treatment is somehow risky for normal tissue. [Cemek et al. \(2006\)](#) showed that *N. sativa* and glutathione treatment significantly antagonize the effects of radiation. Therefore, *N. sativa* may be a beneficial agent in protection against ionizing radiation-related tissue injury. [Assayed \(2010\)](#) investigated the radio-protective potential of *N. sativa* crude oil against hemopoietic adverse effects of gamma irradiation. He found that irradiation resulted in significant reduction in hemolysin antibodies titers and delayed type hypersensitivity reaction of irradiated rats, in addition to significant leukopenia and significant decrease in plasma total protein and globulin concentration and depletion of

lymphoid follicles of spleen and thymus gland. Furthermore, there was a significant increase in malondialdehyde concentration with a significant decrease in plasma GPx, CAT and erythrocyte SOD activities. But oral administration of *N. sativa* oil before irradiation considerably normalized all the above-mentioned criteria; and produced significant regeneration in spleen and thymus lymphoid follicles. Thus *N. sativa* oil is recognized as a promising natural radioprotective agent against immunosuppressive and oxidative effects of ionizing radiation.

### Concluding remarks

The anti-cancer activities of *N. sativa* components were recognized thousands of years ago but proper scientific research with this important traditional medicine is a very recent story. More research works should be emphasized behind this because it is a safe and promising anticancer agent. Specially, researchers should investigate the active ingredients more broadly, because, there is very few authentic reports about the chemical composition of seeds or oil of *N. sativa* exist. Also, the exact molecular mechanisms of thymoquinone and other components on different cancers should be investigated with more emphasis because current understandings are mostly unclear. For example, it is reported that *N. sativa* oil can protect cells from radiation, but the molecular mechanisms behind this is not properly understood. Currently, in some parts of the world, there is a renaissance of interest in traditional remedies. Many investigators now believe that traditional medicine is a promising source of new therapeutics against cancer. Extensive research with *N. sativa* may contribute to the discovery of new anticancer strategies.

### References

1. Aggarwal BB, Kunnumakkara AB, Harikumar KB, Tharakan ST, Sung B, Anand P. Potential of spice-derived phytochemicals for cancer prevention. *Planta Med.* 2008;74:1560–1569. [[PubMed](#)] [[Google Scholar](#)]
2. Al-Ali A, Alkhawajah AA, Randhawa MA, Shaikh NA. Oral and intraperitoneal LD50 of thymoquinone, an active principle of *Nigella sativa*, in mice and rats. *J Ayub Med Coll Abbottabad.* 2008;20:252–257. [[PubMed](#)] [[Google Scholar](#)]
3. Albensi BC, Mattson MP. Evidence for the involvement of TNF and NF-kappaB in hippocampal synaptic plasticity. *Synapse.* 2000;35:151–159. [[PubMed](#)] [[Google Scholar](#)]
4. Alenzi FQ, El-Bolkiny Yel-S, Salem ML. Protective effects of *Nigella sativa* oil and thymoquinone against toxicity induced by the anticancer drug cyclophosphamide. *Br J Biomed Sci.* 2010;67:20–28. [[PubMed](#)] [[Google Scholar](#)]
5. Ali BH, Blunden G. Pharmacological and toxicological properties of *Nigella sativa*. *Phytother Res.* 2003;17:299–305. [[PubMed](#)] [[Google Scholar](#)]
6. Assayed ME. Radioprotective effects of black seed (*Nigella sativa*) oil against hemopoietic damage and immunosuppression in gamma-irradiated rats. *Immunopharmacol Immunotoxicol.* 2010;32:284–296. [[PubMed](#)] [[Google Scholar](#)]
7. Awad EM. In vitro decreases of the fibrinolytic potential of cultured human fibrosarcoma cell line, HT1080, by *Nigella sativa* oil. *Phytomedicine.* 2005;12:100–107. [[PubMed](#)] [[Google Scholar](#)]
8. Badary OA, Gamal El-Din AM. Inhibitory effects of thymoquinone against 20-methylcholanthrene-induced fibrosarcoma tumorigenesis. *Cancer Detect Prev.* 2001;25:362–368. [[PubMed](#)] [[Google Scholar](#)]
9. Badary OA, Taha RA, Gamal el-Din AM, Abdel-Wahab MH. Thymoquinone is a potent superoxide anion scavenger. *Drug Chem Toxicol.* 2003;26:87–98. [[PubMed](#)] [[Google Scholar](#)]



10. Banerjee S, Kaseb AO, Wang Z, Kong D, Mohammad M, Padhye S, Sarkar FH, Mohammad RM. Antitumor activity of gemcitabine and oxaliplatin is augmented by thymoquinone in pancreatic cancer. *Cancer Res.* 2009;69:5575–5583. [[PubMed](#)] [[Google Scholar](#)]
11. Barron J, Benghuzzi H, Tucci M. Effects of thymoquinone and selenium on the proliferation of mg 63 cells in tissue culture. *Biomed Sci Instrum.* 2008;44:434–440. [[PubMed](#)] [[Google Scholar](#)]
12. Bayrak O, Bavbek N, Karatas OF, Bayrak R, Catal F, Cimentepe E, Akbas A, Yildirim E, Unal D, Akcay A. *Nigella sativa* protects against ischaemia/reperfusion injury in rat kidneys. *Nephrol Dial Transplant.* 2008;23:2206–2212. [[PubMed](#)] [[Google Scholar](#)]
13. Bourgou S, Ksouri R, Bellila A, Skandrani I, Falleh H, Marzouk B. Phenolic composition and biological activities of Tunisian *Nigella sativa* L. shoots and roots. *C R Biol.* 2008;331:48–55. [[PubMed](#)] [[Google Scholar](#)]
14. Boskabady MH, Mohsenpoor N, Takaloo L. Antiasthmatic effect of *Nigella sativa* in airways of asthmatic patients. *Phytomedicine.* 2010 Epub ahead of print. [[PubMed](#)] [[Google Scholar](#)]
15. Cemek M, Enginar H, Karaca T, Unak P. In vivo radioprotective effects of *Nigella sativa* L oil and reduced glutathione against irradiation-induced oxidative injury and number of peripheral blood lymphocytes in rats. *Photochem Photobiol.* 2006;82:1691–1696. [[PubMed](#)] [[Google Scholar](#)]
16. Chehl N, Chipitsyna G, Gong Q, Yeo CJ, Arafat HA. Anti-inflammatory effects of the *Nigella sativa* seed extract, thymoquinone, in pancreatic cancer cells. *HPB (Oxford)* 2009;11:373–381. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
17. Chen HC, Hu WX, Liu QX, Li WK, Chen FZ, Rao ZZ, Liu XF, Luo YP, Cao YF. Genetic polymorphisms of metabolic enzymes CYP1A1, CYP2D6, GSTM1 and GSTT1 and leukemia susceptibility. *Eur J Cancer Prev.* 2008;17:251–258. [[PubMed](#)] [[Google Scholar](#)]
18. Dahri AH, Chandiol AM, Rahoo AA, Memon RA. Effect of *Nigella sativa* (kalonji) on serum cholesterol of albino rats. *J Ayub Med Coll Abbottabad.* 2005;17:72–74. [[PubMed](#)] [[Google Scholar](#)]
19. Dehkordi FR, Kamkhah AF. Antihypertensive effect of *Nigella sativa* seed extract in patients with mild hypertension. *Fundam Clin Pharmacol.* 2008;22:447–452. [[PubMed](#)] [[Google Scholar](#)]
20. Ebru U, Burak U, Yusuf S, Reyhan B, Arif K, Faruk TH, Emin M, Aydin K, Atilla II, Semsettin S, Kemal E. Cardioprotective effects of *Nigella sativa* oil on cyclosporine A-induced cardiotoxicity in rats. *Basic Clin Pharmacol Toxicol.* 2008;103:574–580. [[PubMed](#)] [[Google Scholar](#)]
21. Effenberger K, Breyer S, Schobert R. Terpene conjugates of the *Nigella sativa* seed-oil constituent thymoquinone with enhanced efficacy in cancer cells. *Chem Biodivers.* 2010;7:129–139. [[PubMed](#)] [[Google Scholar](#)]
22. El-Aziz MA, Hassan HA, Mohamed MH, Meki AR, Abdel-Ghaffar SK, Hussein MR. The biochemical and morphological alterations following administration of melatonin, retinoic acid and *Nigella sativa* in mammary carcinoma: an animal model. *Int J Exp Pathol.* 2005;86:383–396. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
23. El-Mahdy MA, Zhu Q, Wang QE, Wani G, Wani AA. Thymoquinone induces apoptosis through activation of caspase-8 and mitochondrial events in p53-null myeloblastic leukemia HL-60 cells. *Int J Cancer.* 2005;117:409–417. [[PubMed](#)] [[Google Scholar](#)]
24. EI Shenawy NS, Soliman MF, Reyad SI. The effect of antioxidant properties of aqueous garlic extract and *Nigella sativa* as anti-schistosomiasis agents in mice. *Rev Inst Med Trop Sao Paulo.* 2008;50:29–36. [[PubMed](#)] [[Google Scholar](#)]

25. Farah IO, Begum RA. Effect of *Nigella sativa* (N. sativa L.) and oxidative stress on the survival pattern of MCF-7 breast cancer cells. *Biomed Sci Instrum.* 2003;39:359–364. [[PubMed](#)] [[Google Scholar](#)]
26. Gali-Muhtasib H, Diab-Assaf M, Boltze C, Al-Hmaira J, Hartig R, Roessner A, Schneider-Stock R. Thymoquinone extracted from black seed triggers apoptotic cell death in human colorectal cancer cells via a p53-dependent mechanism. *Int J Oncol.* 2004;25:857–866. [[PubMed](#)] [[Google Scholar](#)]
27. Ghosheh OA, Houdi AA, Crooks PA. High performance liquid chromatographic analysis of the pharmacologically active quinones and related compounds in the oil of the black seed (*Nigella sativa* L.) *J Pharm Biomed Anal.* 1999;19:757–762. [[PubMed](#)] [[Google Scholar](#)]
28. Grundy SM. Recent nutrition research: implications for foods of the future. *Ann Med.* 1991;23:187–193. [[PubMed](#)] [[Google Scholar](#)]
29. Ibrahim ZS, Ishizuka M, Soliman M, ElBohi K, Sobhy W, Muzandu K, Elkattawy AM, Sakamoto KQ, Fujita S. Protection by *Nigella sativa* against carbon tetrachloride-induced downregulation of hepatic cytochrome P450 isozymes in rats. *Jpn J Vet Res.* 2008;56:119–128. [[PubMed](#)] [[Google Scholar](#)]
30. Ismail M, Al-Naqeep G, Chan KW. *Nigella sativa* thymoquinone-rich fraction greatly improves plasma antioxidant capacity and expression of antioxidant genes in hypercholesterolemic rats. *Free Radic Biol Med.* 2010;48:664–672. [[PubMed](#)] [[Google Scholar](#)]
31. Kanter M, Akpolat M, Aktas C. Protective effects of the volatile oil of *Nigella sativa* seeds on beta-cell damage in streptozotocin-induced diabetic rats: a light and electron microscopic study. *J Mol Histol.* 2009;40:379–385. [[PubMed](#)] [[Google Scholar](#)]
32. Kaseb AO, Chinnakannu K, Chen D, Sivanandam A, Tejwani S, Menon M, Dou QP, Reddy GP. Androgen receptor and E2F-1 targeted thymoquinone therapy for hormone-refractory prostate cancer. *Cancer Res.* 2007;67:7782–7788. [[PubMed](#)] [[Google Scholar](#)]
33. Khader M, Bresgen N, Eckl PM. Antimutagenic effects of ethanolic extracts from selected Palestinian medicinal plants. *J Ethnopharmacol.* 2010;127:319–324. [[PubMed](#)] [[Google Scholar](#)]
34. Khan MA, Tania M, Zhang DZ, Chen HC. Antioxidant enzymes and cancer. *Chin J Cancer Res.* 2010;22:87–92. [[Google Scholar](#)]
35. Khan N, Sultana S. Inhibition of two stage renal carcinogenesis, oxidative damage and hyperproliferative response by *Nigella sativa*. *Eur J Cancer Prev.* 2005;14:159–168. [[PubMed](#)] [[Google Scholar](#)]
36. Khattab MM, Nagi MN. Thymoquinone supplementation attenuates hypertension and renal damage in nitric oxide deficient hypertensive rats. *Phytother Res.* 2007;21:410–414. [[PubMed](#)] [[Google Scholar](#)]
37. Mabrouk GM, Moselhy SS, Zohny SF, Ali EM, Helal TE, Amin AA, Khalifa AA. Inhibition of methylnitrosourea (MNU) induced oxidative stress and carcinogenesis by orally administered bee honey and *Nigella* grains in Sprague Dawely rats. *J Exp Clin Cancer Res.* 2002;21:341–346. [[PubMed](#)] [[Google Scholar](#)]
38. Nader MA, el-Agamy DS, Suddek GM. Protective effects of propolis and thymoquinone on development of atherosclerosis in cholesterol-fed rabbits. *Arch Pharm Res.* 2010;33:637–643. [[PubMed](#)] [[Google Scholar](#)]

39. Nagi MN, Almakki HA. Thymoquinone supplementation induces quinone reductase and glutathione transferase in mice liver: possible role in protection against chemical carcinogenesis and toxicity. *Phytother Res.* 2009;23:1295–1298. [[PubMed](#)] [[Google Scholar](#)]
40. *Nigella-sativa-research.com*, author. *Nigella Sativa* - History & Origins. 2010. [February, 03, 2010]. Available at: [http://nigella-sativa-research.com/?page\\_id=41](http://nigella-sativa-research.com/?page_id=41).
41. Norwood AA, Tan M, May M, Tucci M, Benghuzzi H. Comparison of potential chemotherapeutic agents, 5fluoruracil, green tea, and thymoquinone on colon cancer cells. *Biomed Sci Instrum.* 2006;42:350–356. [[PubMed](#)] [[Google Scholar](#)]
42. Oberg F, Haseeb A, Ahnfelt M, Pontén F, Westermarck B, El-Obeid A. Herbal melanin activates TLR4/NF-kappaB signaling pathway. *Phytomedicine.* 2009;16:477–484. [[PubMed](#)] [[Google Scholar](#)]
43. Pari L, Sankaranarayanan C. Beneficial effects of thymoquinone on hepatic key enzymes in streptozotocin-nicotinamide induced diabetic rats. *Life Sci.* 2009;85:830–834. [[PubMed](#)] [[Google Scholar](#)]
44. Ragheb A, Attia A, Eldin WS, Elbarbry F, Gazarin S, Shoker A. The protective effect of thymoquinone, an antioxidant and anti-inflammatory agent, against renal injury: a review. *Saudi J Kidney Dis Transpl.* 2009;20:741–752. [[PubMed](#)] [[Google Scholar](#)]
45. Rooney S, Ryan MF. Modes of action of alpha-hederin and thymoquinone, active constituents of *Nigella sativa*, against HEp-2 cancer cells. *Anticancer Res.* 2005;25:4255–4259. [[PubMed](#)] [[Google Scholar](#)]
46. Salim EI, Fukushima S. Chemopreventive potential of volatile oil from black cumin (*Nigella sativa* L.) seeds against rat colon carcinogenesis. *Nutr Cancer.* 2003;45:195–202. [[PubMed](#)] [[Google Scholar](#)]
47. Salomi MJ, Nair SC, Panikkar KR. Inhibitory effects of *Nigella sativa* and saffron (*Crocus sativus*) on chemical carcinogenesis in mice. *Nutr Cancer.* 1991;16:67–72. [[PubMed](#)] [[Google Scholar](#)]
48. Sethi G, Ahn KS, Aggarwal BB. Targeting nuclear factor-kappa B activation pathway by thymoquinone: role in suppression of antiapoptotic gene products and enhancement of apoptosis. *Mol Cancer Res.* 2008;6:1059–1070. [[PubMed](#)] [[Google Scholar](#)]
49. Shafi G, Munshi A, Hasan TN, Alshatwi AA, Jyothy A, Lei DK. Induction of apoptosis in HeLa cells by chloroform fraction of seed extracts of *Nigella sativa*. *Cancer Cell Int.* 2009;9:29–36. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
50. Sim SC, Ingelman-Sundberg M. The human cytochrome P450 Allele Nomenclature Committee Web site: submission criteria, procedures, and objectives. *Methods Mol Biol.* 2006;320:183–191. [[PubMed](#)] [[Google Scholar](#)]
51. Swamy SM, Huat BT. Intracellular glutathione depletion and reactive oxygen species generation are important in alpha-hederin-induced apoptosis of P388 cells. *Mol Cell Biochem.* 2003;24:127–139. [[PubMed](#)] [[Google Scholar](#)]
52. Thabrew MI, Mitry RR, Morsy MA, Hughes RD. Cytotoxic effects of a decoction of *Nigella sativa*, *Hemidesmus indicus* and *Smilax glabra* on human hepatoma HepG2 cells. *Life Sci.* 2005;77:1319–1330. [[PubMed](#)] [[Google Scholar](#)]
53. Torres MP, Ponnusamy MP, Chakraborty S, Smith LM, Das S, Arafat HA, Batra SK. Effects of Thymoquinone in the Expression of Mucin 4 in Pancreatic Cancer Cells: Implications for the Development of Novel Cancer Therapies. *Mol Cancer Ther.* 2010;9:1419–1431. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

54. Uz E, Bayrak O, Uz E, Kaya A, Bayrak R, Uz B, Turgut FH, Bavbek N, Kanbay M, Akcay A. *Nigella sativa* oil for prevention of chronic cyclosporine nephrotoxicity: an experimental model. *Am J Nephrol*. 2008;28:517–522. [[PubMed](#)] [[Google Scholar](#)]
55. Wikipedia: The free encyclopedia, author. Nigella Sativa. 2010. [April, 03, 2010]. Available at: [http://en.wikipedia.org/wiki/Nigella\\_sativa](http://en.wikipedia.org/wiki/Nigella_sativa).
56. Xuan NT, Shumilina E, Qadri SM, Götz F, Lang F. Effect of thymoquinone on mouse dendritic cells. *Cell Physiol Biochem*. 2010;25:307–314. [[PubMed](#)] [[Google Scholar](#)]
57. Yi T, Cho SG, Yi Z, Pang X, Rodriguez M, Wang Y, Sethi G, Aggarwal BB, Liu M. Thymoquinone inhibits tumor angiogenesis and tumor growth through suppressing AKT and extracellular signal-regulated kinase signaling pathways. *Mol Cancer Ther*. 2008;7:1789–1796. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

---

Articles from African Journal of Traditional, Complementary, and Alternative Medicines are provided here  
courtesy of **African Traditional Herbal Medicine Supporters Initiative**